arXiv:1204.5939v4 [math.DS] 12 Jun 2012

Abstract

Let G be a locally compact group. A random closed subgroup with conjugation-invariant law is called an *invariant random subgroup* or IRS for short. We show that each nonabelian free group has a large “zoo” of IRS’s. This contrasts with results of Stuck-Zimmer which show that there are no non-obvious IRS’s of higher rank semisimple Lie groups with property (T).

Invariant random subgroups of the free group

Lewis Bowen[*]
Texas A&M University

March 3, 2013

Keywords: Invariant random subgroups, ergodic equivalence relations
MSC:37A15, 37A20

Contents

1 Introduction 2

2 Totally Non-Free Actions 5

3 The simplex of invariant measures 8

4 Generic graphings 11

5 Encoding via sub-actions 16

1 Introduction

Let G be a locally compact group. A random closed subgroup with conjugation-invariant law is called an *invariant random subgroup* or IRS for short. These objects arise naturally from the study of group actions. Indeed, suppose G acts on a space X. For $x \in X$, let $\mathrm{Stab}(x) = \{g \in G : gx = x\}$. If μ is a G-invariant probability measure on X and $x \in X$ is chosen at random with law μ then $\mathrm{Stab}(x)$ is an IRS. By [AGV12, Proposition 13] every IRS occurs from this type of construction.

There has been a recent increase in studies of the action of G on its space of subgroups [AGV12, Vo12, ABBGNRS11, Ve11, Sa11, Gr11, Ve10, Bo10, BS06, DS02, GS99, SZ94]. Perhaps the deepest result in the subject is the Stuck-Zimmer Theorem [SZ94] that there are no non-obvious IRS's of higher rank simple Lie groups. More precisely, every ergodic IRS is induced from a lattice subgroup. Another nice result in this area is a complete classification

---

[*] email:lpbowen@math.tamu.edu

IRS's of the infinite symmetric group due to A. Vershik [Ve11]. Our main goal is to show that, by contrast, there is a large zoo of IRS's of any finitely generated nonabelian free group G.

There are three main results of this paper, (1) that the space of ergodic aperiodic IRS's of the free group is homeomorphic to Hilbert space, (2) that every ergodic aperiodic pmp equivalence relation of cost $< r$ is represented by the action of $\mathbb{F}_r$ on its space of subgroups, (3) that every symbolic action of $\mathbb{F}_r$ can be encoded as a sub-action of $\mathbb{F}_r$ on its space of subgroups.

To be precise, we need to introduce some notation. Let $S(G)$ denote the space of all closed subgroups of a locally compact group G with the topology of uniform convergence on compact sets. G acts on $S(G)$ by $g \cdot H := gHg^{-1}$ and $M(G)$ denotes the space of all conjugation-invariant Borel probability measures on $S(G)$. We always consider $M(G)$ with the weak* topology.

Problem 1. Describe $M(G)$ up to affine isomorphism.

Recall that convex closed metrizable subset K of a locally convex linear space is a *simplex* if each point in K is the barycenter of a unique probability measure supported on the subset $\partial_e K$ of extreme points of K. In this case, K is called a *Poulsen simplex* if $\partial_e K$ is dense in K. It is called a *Bauer simplex* if $\partial_e K$ is closed.

For example, an old result states that the space of all shift-invariant Borel probability measures on $\{0,1\}^{\mathbb{Z}}$ is a Poulsen simplex (we will not need this fact). It is known from [LOS78] that there is a unique Poulsen simplex up to affine isomorphism. Moreover, its set of extreme points is homeomorphic to the Hilbert space l^2. On the other hand, there are uncountably many non-isomorphic Bauer simplices. For example, let X be any compact metrizable space. Then the space $P(X)$ of all Borel probability measures on X is a Bauer simplex with $\partial_e P(X)$ homeomorphic to X.

Let $M_e(G) \subset M(G)$ denote the ergodic measures and $M_{fi}(G) \subset M_e(G)$ those measures which are supported on the conjugacy class of a single finite-index subgroup. If G is finitely generated, then each of its finite-index subgroups are finitely generated and because of this, each finite-index subgroup is isolated in the space of subgroups. This implies each element of $M_{fi}(G)$ is isolated in $M_e(G)$. In particular, $M_e(G)$ cannot be connected if G has proper finite-index subgroups.

Let $M_{ie}(G) := M_e(G) \setminus M_{fi}(G)$ and $M_i(G)$ the closed convex hull of $M_{ie}(G)$. Our first main result is:

Theorem 3.1. *If G is a nonabelian free group, then $M_i(G)$ is a Poulsen simplex. In particular, $M_{ie}(G)$ is a dense G_δ subset of $M_i(G)$ and $M_{ie}(G)$ is homeomorphic to the Hilbert space l^2.*

Remark 1. In the course of proving this result, we show that the set of measures $\mu \in M_i(G)$ such that $G \curvearrowright (S(G), \mu)$ is totally non-free is residual in $M_i(G)$. The study of totally non-free actions was introduced by A. Vershik [Ve10].

Remark 2. In order to discuss the case of locally compact groups, let $M_c(G) \subset M_e(G)$ be the set of measures which are supported on the conjugacy class of a single cocompact subgroup. Let $M_{nce} = M_e(G) \setminus M_c(G)$ and $M_{nc}(G)$ be the closed convex hull of $M_{nce}(G)$. Using the ideas of the proof of Theorem 3.1, it can be shown that $M_{nc}(PSL_2(\mathbb{R}))$ is also a Poulsen simplex. By constract, the main results of [GW97] imply that if G has property (T) then $M_{nc}(G)$ and $M(G)$ are Bauer simplices.

Remark 3. Grigorchuk's *space of r-generated marked groups* [Gr84] is (canonically isomorphic with) the space of normal subgroups of the free group of rank r with the topology of uniform convergence on compact subsets (also known as the Chabauty topology). It can be viewed as a subspace of $M_e(\mathbb{F}_r)$ (where $\mathbb{F}_r$ is the free group of rank r). Namely, for each normal subgroup N, let δ_N be the Dirac probability measure supported on $\{N\}$. Then $\{\delta_N : N \lhd \mathbb{F}_r\} \subset M_e(\mathbb{F}_r)$ is a copy of the space of r-generated marked groups. It is known that the space of marked groups contains interesting isolated points [CGP07]. In particular, there are infinite index normal subgroups which are isolated in the space of all normal subgroups. However, these points are *not* isolated in $M_e(\mathbb{F}_r)$ because $M_{ie}(\mathbb{F}_r)$ is pathwise connected by Theorem 3.1.

To explain the next result, we need to recall some notions from the theory of measured equivalence relations. So let (X, μ) be a standard Borel probability space, $E \subset X \times X$ a discrete Borel equivalence relation such that μ is E-invariant (i.e., if $\phi : X \to X$ is a Borel automorphism whose graph is contained in E then $\phi_*\mu = \mu$). We refer to the triple (X, μ, E) as a *discrete pmp equivalence relation.* Two discrete measured equivalence relations (X_i, μ_i, E_i) (for $i = 1, 2$) are *isomorphic* if there exist conull sets $X_i' \subset X_i$ and a measure-space isomorphism $\phi : (X_1', \mu_1) \to (X_2', \mu_2)$ such that $(x, y) \in E_1 \Leftrightarrow (\phi(x), \phi(y)) \in E_2$. More precisely, we only require that ϕ is defined on a set of full measure.

Let E_G denote the orbit equivalence relation on $S(G)$: $E_G := \{(K, gKg^{-1}) : K \in S(G), g \in G\}$. Our second main result is:

Corollary 4.5. *If G is a free group of rank r and (X, μ, E) is an ergodic aperiodic discrete pmp equivalence relation with cost(E) $< r$ then there exists an invariant measure $\lambda \in M(G)$ such that $(S(G), \lambda, E_G)$ is isomorphic to (X, μ, E).*

Remark 4. The authors of [IKT09] and I. Epstein proved that if G is any nonamenable group then it is impossible to classify up to countable structures the free mixing probability measure-preserving actions of G up to orbit-equivalence. Note that if G has a generating set with less than r generators then all of its orbit equivalence relations have cost $< r$.

Remark 5. A well-known result of [FM77] states that any pmp equivalence relation (X, μ, E) can be realized as the orbit-equivalence relation for the action of a countable group G. In other words, given (X, μ, E) there is a countable group G and a measure-preserving action $G \curvearrowright (X, \mu)$ such that $(x, y) \in E \Leftrightarrow \exists g \in G$ such that $gx = y$. The authors then asked whether there exists a group G and an *essentially free* action $G \curvearrowright (X, \mu)$ generating E. The negative answer to this question was obtained by Furman [Fu99]. By contrast, Corollary 4.5 implies that if $\text{cost}(E) < r$ then there is a *totally non-free* action of $\mathbb{F}_r$, the free group of rank r, generating the equivalence relation.

Finally, we provide a general method of constructing invariant subspaces and measures on $S(\mathbb{F}_r)$ from symbolic actions of $\mathbb{F}_r$ (where $\mathbb{F}_r$ is the rank r free group):

Theorem 5.1. *Let $\mathbb{F}_r = \langle s_1, \ldots, s_r \rangle$ be the free group of rank $r \geq 2$, $\mathbb{F}_r' < \mathbb{F}_r$ be the (infinite-index) subgroup generated by $\langle s_1^2, s_2, \ldots, s_r \rangle$, K be a finite set, $\mathbb{F}_r \curvearrowright K^{\mathbb{F}_r}$ be the usual action*

$gx(f) := x(g^{-1}f)$. Let $X \subset K^{\mathbb{F}_r}$ be a closed invariant subspace. Then there exist subspaces $Z \subset Y \subset S(\mathbb{F}_r)$ such that

1. *Z is $\mathbb{F}_r'$-invariant, Y is $\mathbb{F}_r$-invariant,*
2. *there is a finite set $L \subset \mathbb{F}_r$ (depending only on K) such that $Y = \bigcup_{f \in L} fZ$,*
3. *the action $\mathbb{F}_r \curvearrowright X$ is topologically conjugate to the conjugation-action $\mathbb{F}_r' \curvearrowright Z$. More precisely, if $\phi : \mathbb{F}_r \to \mathbb{F}_r'$ is the isomorphism determined by $\phi(s_1) = s_1^2, \phi(s_i) = s_i$ for $2 \le i \le r$ then there is a homeomorphism $\Psi : X \to Z$ such that $\Psi(fx) = \phi(f) \cdot \Psi(x)$ for all $x \in X$ and $f \in \mathbb{F}_r$.*
4. *If η is an $\mathbb{F}_r$-invariant Borel measure on X then there exists an $\mathbb{F}_r$-invariant Borel measure λ on Y such that $\Psi_*\eta = \lambda|Z$ (in particular, $\lambda(Z) > 0$). Moreover, if η is finite then λ is also finite.*

Acknowledgements. Thanks to Miklos Abert, Yair Glasner, Eli Glasner and Anatoly Vershik for stimulating questions on this topic and to Rostislav Grigorchuk for helpful comments. Thanks to Benjy Weiss for pointing out errors in a previous version.

2 Totally Non-Free Actions

Let G be a locally compact group and $G \curvearrowright (X, \mathcal{B}, \mu)$ a pmp (probability-measure-preserving) action. For any Borel set $H \subset G$, let $\mathrm{Fix}(H) = \{x \in X : hx = x \ \forall h \in H\}$. A probability measure preserving action $G \curvearrowright (X, \mathcal{B}, \mu)$ is *totally non-free* if the collection $\{\mathrm{Fix}(H) : H$ a Borel subset of $G\} \subset \mathcal{B}$ generates the Borel sigma-algebra $\mathcal{B}$ (up to sets of measure zero). The study of such actions was initiated in [Ve11].

For $K < G$ let $N_G(K) := \{g \in G : gKg^{-1} = K\}$ denote the normalizer of K and $M_n(G)$ be the set of all measures $\eta \in M(G)$ such that $\eta(\{K \in S(G) : N_G(K) = K\}) = 1$.

Lemma 2.1. *For $\mu \in M_n(G)$, the action $G \curvearrowright (S(G), \mu)$ is totally non-free.*

Proof. Observe that because $\mu \in M_n(G)$ for any Borel set $H \subset G$, $\mu(\mathrm{Fix}(H) \Delta \{K \in S(G) : H \subset K\}) = 0$ where Δ denotes symmetric difference. Therefore, for any Borel set $H \subset G$, $\{K \in S(G) : H \subset K\}$ is contained in the sigma-algebra generated by the sets $\{\mathrm{Fix}(H) : H$ a Borel subset of $G\}$ (up to measure zero). This clearly generates the Borel sigma-algebra. □

The next lemma is used in the proof of Theorem 3.1. Recall that $\mathbb{F}_r$ denotes the free group of rank r.

Lemma 2.2. *For any $r \ge 2$, $M_n(\mathbb{F}_r) \cap M_i(\mathbb{F}_r)$ is dense in $M_i(\mathbb{F}_r)$.*

Proof. Let $S = \{s_1, \ldots, s_r\}$ be a free generating set for $\mathbb{F}_r$. Recall that for any $K < \mathbb{F}_r$, the Schreier coset graph of K has vertex set $K\backslash\mathbb{F}_r$ and for every $s \in S$ and $Kg \in K\backslash\mathbb{F}_r$ there is a directed edge (Kg, Kgs) labeled s. In particular, the Schreier coset graph is a directed graph in which every vertex has in-degree and out-degree equal to r. It may have loops or multiple edges. The automorphism group $\mathrm{Aut}(K\backslash\mathbb{F}_r)$ is the group of all graph automorphisms which preserve the labels and directions on edges. It is isomorphic to $K\backslash N_{\mathbb{F}_r}(K)$ via the map $\phi \in \mathrm{Aut}(K\backslash\mathbb{F}_r) \mapsto \phi(K\backslash\mathbb{F}_r)$. In particular, $N_{\mathbb{F}_r}(K) = K$ if and only if $\mathrm{Aut}(K\backslash\mathbb{F}_r)$ is trivial.

Let Ω denote the set of all pairs (K, x) where $K \in S(\mathbb{F}_r)$ and $x : K\backslash\mathbb{F}_r \to \{0, 1\}$ is a map. For $g \in \mathbb{F}_r$, let $T_g(K, x) = (gKg^{-1}, y)$ where $y : gKg^{-1}\backslash\mathbb{F}_r \to \{0, 1\}$ is the map $y(gKg^{-1}f) = x(Kg^{-1}f)$. This defines an action of $\mathbb{F}_r$ on Ω.

For $\eta \in M(\mathbb{F}_r)$ and $p \in [0, 1]$, let η_p be the measure on Ω whose projection to $S(\mathbb{F}_r)$ is η and whose fiber over $K \in S(\mathbb{F}_r)$ is the Bernoulli measure $u_p^{K\backslash\mathbb{F}_r}$ (where u_p is the probability measure on $\{0, 1\}$ given by $u_p(\{1\}) = p$). This measure is preserved under the action $(T_g)_{g\in\mathbb{F}_r}$.

Before providing a formal description of our construction, let us consider the following non-rigorous intuitive picture. Given an element $(K, x) \in \Omega$, we form a new subgroup K_p whose Schreier coset graph is obtained from the Schreier coset graph $K\backslash\mathbb{F}_r$ by 'splitting' every vertex Kg with $x(Kg) = 1$. More precisely, we replace every such vertex with two vertices v_0, v_1. For every edge e labeled by an element $s \in S$ which is directed into Kg, we now consider the same edge as being directed into v_0. If e is an edge labeled by $s \in S$ directed out of Kg then we consider the same edge as now being directed out of v_1. We also put in new edges: for every $s \in S$ there is an edge labeled s directed from v_0 to v_1 (see figure 1). Let K_p denote the subgroup of $\mathbb{F}_r$ whose Schreier coset graph we just constructed. If (K, x) is chosen at random with law η_p (for some $\eta \in M_i(\mathbb{F}_r)$) then K_p is an invariant random subgroup. The law of K_p tends to η as $p \to 0$ and $N_{\mathbb{F}_r}(K_p) = K_p$ almost surely whenever $0 < p < 1$. This construction has a slight flaw - the subgroup K_p corresponds to the stabilizer of a vertex of its Schreier coset graph, but we did not specify which vertex. A more precise dynamical description follows.



Figure 1: The element x is depicted on the left as a function on the vertices of the Schreier coset graph $K\backslash\mathbb{F}_2$. The Schreier coset graph $K_p\backslash G$ is depicted on the right. Black arrows represent s_1 and white arrows represent s_2.

For $((K,x),i) \in \Omega \times \{0,1\}$ and $s \in S$, define $U_s(K,x,i)$ by

- $U_s(K,x,i) = (T_s(K,x),i)$ if $x(K) = 0$ and $i \in \{0,1\}$;
- $U_s(K,x,0) = (K,x,1)$ if $x(K) = 1$;
- $U_s(K,x,1) = (T_s(K,x),0)$ if $x(K) = 1$.

This defines an action of $\mathbb{F}_r$ on $\Omega \times \{0,1\}$. Now let η'_p be the measure on $\Omega \times \{0,1\}$ satisfying

1. $\eta'_p(E \times \{0\}) = \frac{1}{1+p}\eta_p(E)$ for every $E \subset \{(K,x) \in \Omega : \ x(K) = 0\}$;
2. $\eta'_p(E \times \{1\}) = 0$ for every $E \subset \{(K,x) \in \Omega : \ x(K) = 0\}$;
3. $\eta'_p(E \times \{0\}) = \frac{1}{1+p}\eta_p(E)$ for every $E \subset \{(K,x) \in \Omega : \ x(K) = 1\}$;
4. $\eta'_p(E \times \{1\}) = \frac{1}{1+p}\eta_p(E)$ for every $E \subset \{(K,x) \in \Omega : \ x(K) = 1\}$.

It is easy to check that η'_p is invariant under the $\mathbb{F}_r$ action. Finally, let $\mathrm{Stab} : \Omega \times \{0,1\} \to S(\mathbb{F}_r)$ be the stabilizer map $\mathrm{Stab}((K,x),i)) = \{g \in \mathbb{F}_r : \ g((K,x),i) = ((K,x),i)\}$ and $\mu_p := \mathrm{Stab}_*\eta'_p$. It is evident that μ_p varies continuously with p and $\mu_0 = \eta$. Thus $\lim_{p\to 0} \mu_p = \eta$.

Now suppose $\eta \in M_i(\mathbb{F}_r)$. We claim that $\mu_p \in M_n(\mathbb{F}_r) \cap M_i(\mathbb{F}_r)$. It is immediate that $\mu_p \in M_i(\mathbb{F}_r)$ because the Schreier coset graphs of η-a.e. K are "expanded" in the construction of μ_p. If $0 < p < 1$ and L is a random subgroup with law μ_p then $\mathrm{Aut}(L\backslash\mathbb{F}_r)$ is trivial almost surely. Therefore $\mu_p \in M_n(\mathbb{F}_r)$ as required.

□

Lemma 2.3. *The subset $M_n(\mathbb{F}_r)$ is a G_δ-subset of $M(\mathbb{F}_r)$.*

Proof. Given a finite set $F \subset \mathbb{F}_r$ and an element $g \in G$, let $X(F,g)$ be the set of $K \in S(\mathbb{F}_r)$ such that $g(F \cap K)g^{-1} \subset K$. Let $Y(g) = \{K \in S(\mathbb{F}_r) : \ g \notin K\}$. Both $X(F,g)$ and $Y(g)$ are clopen sets. Let $Z(g) = Y(g) \cap \bigcap_F X(F,g)$. Then $Z(g)$ is a closed set. Note that $K \in Z(g)$ if and only if $g \in N_{\mathbb{F}_r}(K) \setminus K$.

Let $O(g,\epsilon)$ be the set of all measures μ in $M(\mathbb{F}_r)$ such that $\mu(Z(g)) < \epsilon$. Then $O(g,\epsilon)$ is open and

$$M_n(\mathbb{F}_r) = \cap_{g\in\mathbb{F}_r} \cap_{m=1}^{\infty} O(g, 1/m).$$

□

Corollary 2.4. *The set of measures $\mu \in M_i(\mathbb{F}_r)$ such that $\mathbb{F}_r \curvearrowright (S(\mathbb{F}_r),\mu)$ is totally non-free is residual in $M_i(\mathbb{F}_r)$.*

3 The simplex of invariant measures

Theorem 3.1. *If G is a nonabelian free group, then $M_i(G)$ is a Poulsen simplex. In particular, $M_{ie}(G)$ is a dense G_δ subset of $M_i(G)$ and $M_{ie}(G)$ is homeomorphic to the Hilbert space l^2.*

Before getting to the proof, we provide here an intuitive but nonrigorous sketch. Let $G = \langle s_1, \dots s_r \rangle$ and $S = \{s_1, \dots, s_r\}$. We will employ the following ‘surgical procedure’ frequently. If for $i = 1, 2$, K_i are two subgroups of G and $K_i g_i \in K_i \backslash G$ are chosen then we can “connect” the Schreier coset graphs $K_i \backslash G$ together at Kg_i, to obtain a new Schreier coset graph $K_3 \backslash G$ by

1. replace the vertex Kg_i with two vertices v_i, w_i;
2. for every edge directed into Kg_i, we form a new edge with the same label and initial vertex going into v_i;
3. for every edge directed out of Kg_i, we form a new edge with the same label and terminal vertex going out of w_i;
4. we add an edge labeled s_1 from v_i to w_i;
5. for every $2 \le i \le r$, we add an edge labeled s_i from v_1 to w_2 and an edge labeled s_i from v_2 to w_1.

See figure 2.



Figure 2: On the left, we see portions of the Schreier coset graphs $K_i \backslash G$ for $i = 1, 2$. On the right, the Schreier coset graph of a new subgroup, obtained from surgery. Black arrows represent s_1 and white arrows represent s_2.

Let $\eta \in M_n(G) \cap M_i(G)$. By Lemma 2.2, it suffices to show that η is a weak* limit of a sequence of measures in $M_{ie}(G)$. Let $0 < p < 1$ and let K be a random subgroup with law η. For every vertex of $K \backslash G$, flip a $(p, 1-p)$-weighted coin. If the coin lands on heads (with probability p), then we choose an independent η-random subgroup and attach its Schreier coset graph to $K \backslash G$ by the surgery procedure above. For each vertex of each new attached graph, we flip a $(p, 1-p)$-weighted coin. If the coin lands on heads (with probability p), then we choose an independent η-random subgroup and attach its Schreier coset graph by the surgery procedure above. We continue this construction for every new attached graph. See figure 3. The result is the Schreier coset graph of an invariant random subgroup K_p. The law of K_p is ergodic and tends to η as $p \to 0$.



Figure 3: The ellipses represent independent copies of $K\backslash G$ where K is η-random. They are attached via the surgery procedure described above at Bernoulli random points.

Proof of Theorem 3.1. It is a well-known general result that the extreme points of a simplex are always a G_δ. To show that it is dense, we define an action of $G = \mathbb{F}_r$ on a space of decorated trees. Let Ω' be the set of all quintuples (T, ϕ, v, K, σ) such that

1. $T = (V(T), E(T))$ is a tree in which every vertex has countable valence;
2. $v \in V(T)$ is a distinguished vertex;
3. ϕ is a map from the half-edges of T to $S(\mathbb{F}_r)$ such that
    - if two half-edges $h_1 \neq h_2$ share a vertex then $\phi(h_1)$ is conjugate to $\phi(h_2)$ and $\phi(h_1) \neq \phi(h_2)$;
4. $K \in S(\mathbb{F}_r)$ is conjugate to $\phi(h)$ where h is an half-edge incident to v;
5. $\sigma \in \{0, 1, 2\}$ and $\sigma = 0$ if and only if $K \neq \phi(h)$ for some half-edge h incident to v.

We consider two quintuples $(T_i, \phi_i, v_i, K_i, \sigma_i) \in \Omega'$ $(i = 1, 2)$ to be isomorphic if there is a graph-isomorphism Ψ from T_1 to T_2 which preserves all the data: $\Psi \circ \phi_2 = \phi_1$, $\Psi(v_1) = v_2, K_1 = K_2, \sigma_1 = \sigma_2$. Let Ω denote the set of isomorphism classes in Ω'. By abuse of notation, we will usually denote elements of Ω as quintuples (T, ϕ, v, K, σ).

We denote half-edges of T by (v, e) where $v \in V(T)$ is the vertex of the half-edge and $e \in E(T)$ is the edge containing it. Let $S = \{s_1, \ldots, s_r\}$ be a free generating set for $\mathbb{F}_r$. We define an action of $\mathbb{F}_r$ on Ω by the following rules:

1. for any $s \in S \cup S^{-1}$, $s(T, \phi, v, K, 0) := (T, \phi, v, sKs^{-1}, 0)$ if sKs^{-1} is not equal to $\phi(v, e)$ for any $e \in E(T)$.

2. for any $s \in S$, $s(T,\phi,v,K,0) := (T,\phi,v,sKs^{-1},1)$ if $sKs^{-1} = \phi(v,e)$ for some $e \in E(T)$.
3. $s_1(T,\phi,v,K,1) := (T,\phi,v,s_1Ks_1^{-1},2)$.
4. for any $2 \le i \le r$, $s_i(T,\phi,v,K,1) := (T,\phi,w,L,2)$ where $\{v,w\} = e$ is the unique edge with $\phi(v,e) = K$ and $L = \phi(w,e)$.

These rules uniquely determine an action of $\mathbb{F}_r$ on Ω.

We will construct a family of invariant measures on Ω from a given measure $\eta \in M_i(\mathbb{F}_r) \cap M_n(\mathbb{F}_r)$. But first we need two other spaces. For $K \in S(\mathbb{F}_r)$, let $[K]$ denote the conjugacy class of K. Let $S'(\mathbb{F}_r)$ be the space of all pairs (K,Q) where $K \in S(\mathbb{F}_r)$ and $Q \subset [K]$. Given $K \in S(\mathbb{F}_r)$ and $p \in [0,1]$, let u_p^K be the probability measure on the space $2^{[K]}$ of all subsets of $[K]$ determined by: for any disjoint finite sets $V, W \subset [K]$,

$$u_p^K(\{W \subset Q \text{ and } V \cap Q = \emptyset\}) = p^{|W|}(1-p)^{|V|}.$$

Let η'_p be the probability measure on $S'(\mathbb{F}_r)$ given by

$$d\eta'_p(K,Q) = du_p^K(Q)d\eta(K).$$

Because $\eta \in M_i(\mathbb{F}_r) \cap M_n(\mathbb{F}_r)$, if $(K,Q) \in S'(\mathbb{F}_r)$ has law η'_p then $|Q| = \infty$ almost surely.

Next let $S''(\mathbb{F}_r)$ be the space of all pairs (K,Q,σ) where $\sigma \in \{0,1,2\}$ satisfies $\sigma = 0$ if and only if $K \notin Q$. Let $(K,Q,\sigma) \in S''(\mathbb{F}_r)$ be the random element satisfying

1. the law of (K,Q) is $\left(\frac{1-p}{1+p}\eta'_p|\{(K,Q) :\ K \notin Q\} + \frac{2p}{1+p}\eta'_p|\{(K,Q) :\ K \in Q\}\right)$ (where $|$ denotes restriction);
2. the law of σ given that $K \in Q$ is uniformly distributed on $\{1,2\}$.

Let η''_p be the law of (K,Q,σ).

Let T denote a tree in which every vertex has countable valence. Let v_0 be a distinguished vertex of T. Let $D_{v_0} = (K_{v_0}, Q_{v_0}, \sigma_{v_0}) \in S''(\mathbb{F}_r)$ be a random element with law η''_p. For each vertex $w \in V(T) \setminus \{v_0\}$, let $D_w = (K_w, Q_w) \in S'(\mathbb{F}_r)$ be a random element with law η''_p conditioned on $K_w \in Q_w$. We require that the variables $\{D_w :\ w \in V(T)\}$ are independent.

Let ϕ be a map on half-edges of T satisfying

$$\forall w \in V(T), \forall K \in Q_w\ \exists! e \in E(T) \text{ such that } w \in e \text{ and } \phi(w,e) = K. \qquad (1)$$

This does not determine the law of ϕ, however we let the law of ϕ be maximally independent subject to this condition. One way to achieve this is as follows. First choose $\phi(v_0,e)$ for all e. It does not matter how we make this choice because all choices are equivalent up to automorphisms of the tree which fix v_0. Second, for each vertex w adjacent to v_0 by an edge e, let $\phi(w,e) = K_w$. Choose $\phi(w,e')$ for all other edges e' incident to such a vertex w in any way subject to the condition above. Again, all choices are equivalent up to automorphisms of the tree which fix v_0, so the actual choice is irrelevant. Assuming, for induction, that ϕ

has been defined for all half edges incident to vertices in the radius n ball centered at v_0. If w is in the radius $n+1$ and e is the unique edge connecting w to the n-ball then let $\phi(w,e) = K_w$ and choose $\phi(w,e')$ for all other edges e' incident to w arbitrarily subject to (1). The actual choice is irrelevant because of tree automorphisms. This determines the law of ϕ.

Finally let $(T, \phi, v_0, K_{v_0}, \sigma_{v_0}) \in \Omega$ be the random element constructed above and let μ_p be its law. So μ_p is a Borel probability measure on Ω. Observe that

1. μ_p is invariant under the $\mathbb{F}_r$-action,
2. μ_p is ergodic (even if η is not!),
3. $\lim_{p\to 0} \mathrm{Stab}_*(\mu_p) = \eta$ where $\mathrm{Stab} : \Omega \to S(\mathbb{F}_r)$ is the map $\mathrm{Stab}(z) = \{g \in \mathbb{F}_r : \ gz = z\}$.

We leave these statements as exercises. To give the reader some indication on how to prove ergodicity, let ψ be an $\mathbb{F}_r$-invariant measurable function on Ω. Because of the independence properties of the construction, it is clear that ψ does not depend on finite radius information (namely, because one can 'move' the root vertex to a place far from its initial place so that its new radius n neighborhood is disjoint from its initial radius n neighborhood). More precisely, if two elements $\tau, \eta \in \Omega$ are such that there is an isomorphism from the tree of τ to the tree of η which preserves all the half-edge labels *outside* the ball of radius n, then $\psi(\tau) = \psi(\eta)$ (excluding a measure zero set of τ, η). Therefore, the $\mathbb{F}_r$-invariant sigma-algebra is contained in the "tail"-sigma-algebra which consists of all measurable sets which do not depend on finite radius information in the sense given above. This tail sigma algebra is trivial because of the independence properties of the construction. This implies ergodicity.

It follows that η is in the closure of $M_{ie}(\mathbb{F}_r)$. Because $\eta \in M_i(\mathbb{F}_r) \cap M_n(\mathbb{F}_r)$ is arbitrary, Lemma 2.2 implies $M_{ie}(\mathbb{F}_r)$ is dense in $M_i(\mathbb{F}_r)$. ☐

4 Generic graphings

Let (X, μ) be a standard Borel probability space and $E \subset X \times X$ be a discrete Borel equivalence relation such that μ is E-invariant. We will usually assume that E is ergodic and aperiodic (which means that a.e. equivalence class is infinite). We use [Ke10] as a general reference.

The full group of E, denoted $[E]$, is the group of all (equivalence classes of) invertible Borel transformations f such that $graph(f) = \{(x, fx) : \ x \in X\} \subset E$. Two transformations are equivalent if they agree on a conull subset. By [Ke10, Proposition 3.2], $[E]$ with the uniform metric, defined by

$$d_u(\phi, \psi) = \mu(\{x \in X : \ \phi(x) \neq \psi(x)\}),$$

is a Polish group.

From now on, we assume that E is aperiodic and ergodic. To set notation, let $\mathrm{Hom}(\mathbb{F}_r, [E])$ denote the set of all homomorphisms from the rank r free group $\mathbb{F}_r = \langle s_1, \ldots, s_r \rangle$ to $[E]$.

We identify $\mathrm{Hom}(\mathbb{F}_r, [E])$ with the product space $[E]^r$ in the obvious way and endow it with the product of uniform topologies. Thus it is a Polish space. For latter purposes it will be useful to have the following explicit metric on $\mathrm{Hom}(\mathbb{F}_r, [E])$:

$$\mathbf{d}_u(\alpha, \beta) := \sum_{i=1}^{r} d_u(\alpha(s_i), \beta(s_i))$$

for $\alpha, \beta \in \mathrm{Hom}(\mathbb{F}_r, [E])$. Also let $E_\beta \subset E$ be the subequivalence relation generated by β. Precisely, $(x, y) \in E_\beta$ if there exists $w \in \mathbb{F}_r$ such that $\beta(w)x = y$.

Definition 1. Let (X, μ, E) be a discrete pmp equivalence relation. Let $[[E]]$ be the set of all Borel isomorphisms $\phi : A \to B$ such that $A, B \subset X$ are Borel and the graph of ϕ is contained in E. A subset $\Gamma \subset [[E]]$ is a *graphing* if for a.e. $x \in X$ and every y with $(x, y) \in E$ there is a sequence $\gamma_1, \ldots, \gamma_n \in \Gamma$ and $\epsilon_1, \ldots, \epsilon_n \in \{-1, 1\}$ such that $\gamma_1^{\epsilon_1} \cdots \gamma_n^{\epsilon_n} x = y$. The *cost* of Γ is defined by

$$\mathrm{cost}(\Gamma) = \sum_{\gamma \in \Gamma} \mu(\mathrm{domain}(\gamma)).$$

The *cost* of E is the infimum of $\mathrm{cost}(\Gamma)$ over all graphings Γ of E.

Lemma 4.1. *Let $Hom_g(\mathbb{F}_r, [E]) \subset Hom(\mathbb{F}_r, [E])$ be the set of all homomorphisms α such that $\{\alpha(s_i) : 1 \le i \le r\}$ is a graphing of E (i.e., $E_\alpha = E$ up to measure zero). If either $r > cost(E)$ or ($r = cost(E)$ and E is treeable) then $Hom_g(\mathbb{F}_r, [E])$ is a non-empty G_δ-subset of $Hom(\mathbb{F}_r, [E])$.*

Proof. By [Hj06, Proposition 1.1], $\mathrm{Hom}_g(\mathbb{F}_r, [E])$ is nonempty. So let $\beta \in \mathrm{Hom}_g(\mathbb{F}_r, [E])$. For $w \in \mathbb{F}_r$ and $\epsilon > 0$, let $O(w, \epsilon)$ be the set of all $\gamma \in \mathrm{Hom}(\mathbb{F}_r, [E])$ such that

$$\mu(\{x \in X : (x, \beta(w)x) \in E_\gamma\}) > \epsilon.$$

This set is open and

$$\mathrm{Hom}_g(\mathbb{F}_r, [E]) = \bigcap_{w \in \mathbb{F}_r} \bigcap_{n=1}^{\infty} O(w, 1 - 1/n).$$

This shows $\mathrm{Hom}_g(\mathbb{F}_r, [E])$ is a G_δ. ☐

An element $f \in [E]$ is *aperiodic* if a.e. f-orbit in X is infinite. For somewhat technical reasons (Theorem 4.4), we will be interested in the set $\mathrm{Hom}'_g(\mathbb{F}_r, [E])$ of all homomorphisms $\alpha \in \mathrm{Hom}_g(\mathbb{F}_r, [E])$ such that $\alpha(s_1)$ is ergodic and aperiodic.

Lemma 4.2. *If E is ergodic aperiodic and $cost(E) < r$ then the set $Hom'_g(\mathbb{F}_r, [E])$ is a nonempty G_δ subset of $Hom(\mathbb{F}_r, [E])$.*

Proof. The fact that $\mathrm{Hom}'_g(\mathbb{F}_r, [E])$ is nonempty can be seen from the proof of [KM04, Lemma 27.7] (just choose φ_1 to be ergodic). That lemma is attributed to Hjorth-Kechris. By Lemma 4.1, $\mathrm{Hom}_g(\mathbb{F}_r, [E])$ is a G_δ. Because E is aperiodic, every ergodic element $f \in [E]$ is aperiodic. So it suffices to show that the set of ergodic elements in $[E]$ is a G_δ subset. This is part of [Ke10, Theorem 3.6]. ☐

We will say that E is *hyperfinite* if there exists a set $Y \subset X$ with $\mu(Y) = 1$ and an increasing sequence $\{R_i\}_{i=1}^\infty$ of finite measurable equivalence relations on Y such that $E \cap (Y \times Y) = \cup_{i=1}^\infty R_i$. We say that a subset $K \subset [[E]]$ *generates* E if E is the smallest equivalence relation containing, for each $k \in K$ the graph $\{(x, kx) : x \in \text{domain}(k)\}$ (up to sets of measure zero).

Lemma 4.3. *Suppose E is ergodic and aperiodic. Let $f \in [E]$. Then there exists an aperiodic Borel hyperfinite subequivalence relation $F \subset E$ such that $f \in [F] < [E]$. Moreover there is an aperiodic $g \in [F]$ which generates F.*

Proof. By abuse of notation, we identify $f : X \to X$ as a Borel map with graph in E (instead of an equivalence class of maps). Let $R \subset E$ be the subequivalence relation generated by f. So $R = \{(x, f^n x) : n \in \mathbb{Z}\}$. For $x \in X$, let $[x]_R$ denote the R-class of x. Let Y be the set of all $x \in X$ such that $[x]_R$ is finite. If $\mu(Y) = 0$ then we may set $F := R$ and $g := f$. So without loss of generality, we assume $\mu(Y) > 0$.

Because Y is a standard Borel probability space, there exists an injective Borel map $\phi : Y \to \mathbb{R}$. Let Z be the set of all $z \in Y$ such that $\phi(z) \le \phi(x)$ for every $x \in [z]_R$. Then Z meets every $R|Y$-equivalence class in exactly one point, where $R|Y$ denotes the restriction of R to Y. In other words, $|[x]_R \cap Z| = 1$ for a.e. $x \in Y$. Because $\mu(Y) > 0$, we must have $\mu(Z) > 0$.

Because E is aperiodic and ergodic, it follows that E restricted to Z is aperiodic and ergodic. By [Ke10, Theorems 3.5, 3.6] there exists an aperiodic ergodic Borel hyperfinite subequivalence relation $G \subset E|Z$. Let F' be the subequivalence relation of $E|Y$ generated by G and $R|Y$. That is, F' is the smallest equivalence relation on Y with $G \cup R|Y \subset F'$.

We claim that F' is hyperfinite. Because G is hyperfinite, there exists an increasing sequence $\{G_i\}_{i=1}^\infty$ of finite subequivalence relations $G_i \subset G$ such that $G = \cup_{i=1}^\infty G_i$ (ignoring a measure zero set). Let G'_i be the subequivalence relation of E generated by G_i and $R|Y$. These are finite subequivalence relations. Indeed, if $x \in X$ and z is the unique element in $|[x]_R \cap Z|$ then the G'_i equivalence class of x is $\cup\{[y]_R : (y, z) \in G_i\}$. By construction, $\cup_{i=1}^\infty G'_i = F'$ (ignoring a measure zero set).

We claim that F' is ergodic. To see this let $W \subset Y$ be a measurable set of positive measure equal to a union of F'-classes. Because $|[x]_R \cap Z| = 1$ for a.e. $x \in Y$, it follows that $W \cap Z$ is a positive measure set. Because G is ergodic, $Z \subset W$ (up to a set of measure zero). Because every $R|Y$ equivalence class intersects Z nontrivially, $W = Y$ up to a set of measure zero. Because W is arbitrary, F' is ergodic.

Let $F = F' \cup R$. Because F' is the disjoint union of F' and $R|(X \setminus Y)$, it is an aperiodic Borel hyperfinite subequivalence relation of E with $f \in [F]$ (because $R \subset F$). Because F' is ergodic, aperiodic and hyperfinite there exists an aperiodic element $f' \in [F']$ which generates F'. So define $g \in [F]$ by

$$gx = \begin{cases} fx & \text{if } x \in X \setminus Y \\ f'x & \text{if } x \in Y \end{cases}$$

By construction, g generates F. □

For $\beta \in \mathrm{Hom}(\mathbb{F}_r, [E])$ let $Stab^\beta : X \to S(\mathbb{F}_r)$ be the stabilizer map: $Stab^\beta(x) = \{g \in \mathbb{F}_r : \beta(g)x = x\}$. Observe that for any $g \in \mathbb{F}_r$ and $x \in X$, $Stab^\beta(\beta(g)x) = gStab^\beta(x)g^{-1}$. In other words, Stab^β is $\mathbb{F}_r$-equivariant. Let $\mu_\beta \in M(\mathbb{F}_r)$ be the pushforward measure $\mu_\beta := \mathrm{Stab}^\beta_* \mu$. Therefore, Stab^β defines a factor map from $\mathbb{F}_r \curvearrowright^\beta (X, \mu)$ to $\mathbb{F}_r \curvearrowright (S(\mathbb{F}_r), \mu_\beta)$.

Theorem 4.4. *Let (X, μ, E) be an ergodic aperiodic discrete pmp equivalence relation. Let $Hom_{isom}(\mathbb{F}_r, [E])$ be the set of homomorphisms $\beta \in Hom(\mathbb{F}_r, [E])$ such that $Stab^\beta$ is an isomorphism from (X, μ) to $(S(\mathbb{F}_r), \mu_\beta)$. If $r > cost(E)$ then $Hom_{isom}(\mathbb{F}_r, [E]) \cap Hom'_g(\mathbb{F}_r, [E])$ is a dense G_δ subset of $Hom'_g(\mathbb{F}_r, [E])$. In particular, it is nonempty.*

Proof. Let $\{\xi_i\}_{i=1}^\infty$ be an increasing sequence of finite Borel partitions of X such that $\bigvee_{i=1}^\infty \xi_i$ is the partition into points; i.e. for every $x \neq y \in X$ there exists i such that x and y are in different partition elements of ξ_i.

For $W \subset \mathbb{F}_r$, let $\pi_W : S(\mathbb{F}_r) \to 2^W$ be the projection map $\pi_W(H) = H \cap W$ (where 2^W denote the set of all subsets of W). For every $n \in \mathbb{N}, \epsilon > 0$ and finite $W \subset \mathbb{F}_r$, let $O(W, n, \epsilon)$ be the set of all $\beta \in \mathrm{Hom}(\mathbb{F}_r, [E])$ such that for every $A, B \in \xi_n$ with $A \neq B$ there exist sets $A' \subset A, B' \subset B$ with

- $\mu(A \setminus A') + \mu(B \setminus B') < \epsilon$ and
- $\pi_W(\mathrm{Stab}^\beta(A')) \cap \pi_W(\mathrm{Stab}^\beta(B')) = \emptyset$.

Each $O(W, n, \epsilon)$ is open in $\mathrm{Hom}(\mathbb{F}_r, [E])$ because for any $\beta \in \mathrm{Hom}(\mathbb{F}_r, [E])$ the set of $\beta' \in \mathrm{Hom}(\mathbb{F}_r, [E])$ which agrees with β on W except for a set of measure $< \epsilon$ is open. Also observe that if $W \subset V$ then $O(W, n, \epsilon) \subset O(V, n, \epsilon)$. Thus, $O(n, \epsilon) := \cup_{W \subset \mathbb{F}_r} O(W, n, \epsilon)$ is open.

We claim that

$$\mathrm{Hom}_{isom}(\mathbb{F}_r, [E]) = \cap_{n=1}^\infty \cap_{k=1}^\infty O(n, 1/k).$$

The inclusion $\subset$ is obvious. To see the other direction, let $\beta \in \cap_{n=1}^\infty \cap_{k=1}^\infty O(n, 1/k)$ and $\mu_\beta = \mathrm{Stab}^\beta_* \mu$. Observe that $\mu_\beta(\mathrm{Stab}^\beta(A) \cap \mathrm{Stab}^\beta(B)) = 0$ for any $A \neq B$ in ξ_n for any n. Since

$$\mu_\beta(\mathrm{Stab}^\beta(A)) = \mu((\mathrm{Stab}^\beta)^{-1}(\mathrm{Stab}^\beta(A)))$$

this implies $\mu_\beta(\mathrm{Stab}^\beta(A)) = \mu(A)$ for every $A \in \xi_n$ for any n. Therefore β induces a measure-algebra isomorphism from the measure-algebra of μ to the measure-algebra of μ_β. This implies that β is a measure-space isomorphism (see e.g. [Zi84, Corollary B.7]). So $\beta \in \mathrm{Hom}_{isom}(\mathbb{F}_r, [E])$ which proves the equality above. So $\mathrm{Hom}_{isom}(\mathbb{F}_r, [E])$ is a G_δ.

Let $\alpha \in \mathrm{Hom}'_g(\mathbb{F}_r, [E])$. Let $\mu_\alpha = (\mathrm{Stab}^\alpha)_* \mu \in M(\mathbb{F}_r)$ be the measure induced by α. Because $\mathrm{cost}(E) < r$, and $\{\alpha(s_i)\}_{i=1}^r$ is a graphing, it follows that the cost of the orbit equivalence relation $\mathbb{F}_r \curvearrowright (S(\mathbb{F}_r), \mu_\alpha)$ is also less than r. Therefore, there exists a shortest element $w \in \mathbb{F}_r \setminus \{e\}$ such that $\mu_\alpha(\{H \in S(\mathbb{F}_r) : w \in H\}) > 0$. Observe that every cyclic conjugate of w also satisfies this property. Because $\alpha(s_1)$ is aperiodic, the reduced form of w must contain a generator other than s_1. So after replacing w with one of its cyclic conjugates, reordering $s_2, \ldots, s_r$ if necessary, and replacing s_r with s_r^{-1} if necessary, we may assume that the last element in the reduced form of w is s_r.

Let $S_w = \{H \in S(\mathbb{F}_r) : \ w \in H\}$. We will construct a positive measure subset $S'_w \subset S_w$ such that $\{s_1, \ldots, s_{r-1}, s_r|(S(\mathbb{F}_r) \setminus S'_w)\}$ is a graphing of the orbit-equivalence relation of $\mathbb{F}_r \curvearrowright S(\mathbb{F}_r)$ with respect to μ_α (where $s_r|(S(\mathbb{F}_r) \setminus S'_w)$ denotes the restriction of s_r to $S(\mathbb{F}_r) \setminus S'_w$). To do this, let $w = t_1 \cdots t_n$ be the reduced form of w (so $t_n = s_r$ for example). For $H \in S_w$, let $C_w(H) = \{(t_1 \cdots t_i)^{-1} H (t_1 \cdots t_i) : \ 1 \le i < n\} \subset S(\mathbb{F}_r)$. Observe that because w is a shortest word such that $\mu_\alpha(S_w) > 0$, we must have that $H \notin C_w(H)$ (for a.e. $H \in S_w$).

Let F be the set of all pairs $(H, H') \in S_w \times S_w$ such that either $H \in C_n(H')$ or $H' \in C_n(H)$. We interpret F to be the set of edges in a graph with vertex set S_w. Observe that every vertex of this graph has degree at most n^2. By [KST99, Proposition 4.6], there exists a proper coloring of this graph with only a finite set of colors. This means that there is a measurable map $\phi : S_w \to \Omega$ (where Ω is a finite set) such that for every $(H, H') \in F$ with $H \ne H'$, $\phi(H) \ne \phi(H')$.

Let $\omega \in \Omega$ be an element such that $\mu_\alpha(\phi^{-1}(\omega)) > 0$. Let $S'_w = \phi^{-1}(\omega)$. We claim that $\{s_1, \ldots, s_{r-1}, s_r|(S(\mathbb{F}_r) \setminus S'_w)\}$ is a graphing of the orbit-equivalence relation of $\mathbb{F}_r \curvearrowright S(\mathbb{F}_r)$ with respect to μ_α. It suffices to show that if $H \in S'_w$ then $(H, s_r H s_r^{-1})$ is in the equivalence relation generated by $\{s_1, \ldots, s_{r-1}, s_r|(S(\mathbb{F}_r) \setminus S'_w)\}$. Because $H \in S_w$, $wHw^{-1} = H$. Because $s_r = t_n$, $s_r H s_r^{-1} = (t_1 \cdots t_{n-1})^{-1} H (t_1 \cdots t_{n-1})$. By choice of S'_w, $C_w(H) \cap S'_w = \emptyset$. So there is a path $H, t_1^{-1} H t_1, (t_1 t_2)^{-1} H (t_1 t_2), \ldots, (t_1 \cdots t_{n-1})^{-1} H (t_1 \cdots t_{n-1})$ contained in the graph determined by $\{s_1, \ldots, s_{r-1}, s_r|(S(\mathbb{F}_r) \setminus S'_w)\}$ from H to $s_r H s_r^{-1}$ as required.

Let $\epsilon > 0$. By choosing S'_w smaller if necessary we may assume $0 < \mu_\alpha(S'_w) < \epsilon$. Let $B := (\mathrm{Stab}^\alpha)^{-1}(S'_w) \subset X$. Then $\Phi := \{\alpha(s_1), \ldots, \alpha(s_{r-1}), \alpha(s_r)|_{X \setminus B}\}$ is a graphing of E.

Define a map ψ_0 on $X \setminus B$ by $\psi_0(x) = \alpha(s_r)^n x$ where n is the smallest positive integer such that $\alpha(s_r)^n x \in X \setminus B$. Such an integer exists for a.e. $x \in X \setminus B$ by Poincaré's Recurrence Theorem. For every positive integer k let C_k be the set of all $x \in X \setminus B$ such that $\psi_0^k(x) = x$ and $k \ge 1$ is the smallest number with this property.

There exists an integer $N > 0$ such that $\mu(\cup_{i=N}^\infty C_i) < \epsilon$. Suppose that $\mu(\cup_{i=N}^\infty C_i) > 0$. By Lemma 4.3, there exists an aperiodic subequivalence relation $F \subset E|(\cup_{i=N}^\infty C_i)$ such that $\psi_0|(\cup_{i=N}^\infty C_i) \in [F]$. Moreover, there is an aperiodic element $f \in [F]$ which generates F. Now define ψ on $X \setminus B$ by $\psi(x) = \psi_0(x)$ is $x \notin \cup_{i=N}^\infty C_i$ and $\psi(x) = f(x)$ if $x \in \cup_{i=N}^\infty C_i$. Note: the equivalence relation generated by ψ contains the graph of $\alpha(s_r)|(X \setminus B)$.

Let $\alpha(s_1)_B : B \to B$ be the first return time map $\alpha(s_1)_B x = \alpha(s_1)^n x$ where n is the smallest positive integer such that $\alpha(s_1)^n x \in B$. Because $\alpha(s_1)$ is ergodic the action of $\alpha(s_1)_B$ is also ergodic. By Rohlin's Theorem, there exists a countable Borel partition γ of B such that if $\mathcal{B}_B$ is the smallest $\alpha(s_1)_B$-invariant sigma-algebra containing γ, then $\mathcal{B}_B$ is the sigma-algebra of all measurable sets (up to sets of measure zero) of B.

Let $\gamma = \{P_i\}_{i=1}^\infty$ and choose elements $f_i \in [E|P_i]$ (the full group of E restricted to P_i) such that f_i has period $i + N$. To be precise, we require that for a.e. $x \in P_i$, $f_i^{i+N} x = x$ and if $1 \le j < i + N$ then $f_i^j x \ne x$. Such an element f_i exists by [Ke10, Theorem 3.3].

We now extend the definition of ψ to B by setting $\psi(x) = f_i(x)$ if $x \in P_i$. Also define a homomorphism $\beta : \mathbb{F}_r \to [E]$ by $\beta(s_i) = \alpha(s_i)$ if $1 \le i < r$ and $\beta(s_r) = \psi$. We claim that

1. $d_u(\alpha, \beta) := \sum_{i=1}^r d_u(\alpha(s_i), \beta(s_i)) \le 2\epsilon$;
2. $\beta \in \mathrm{Hom}'_g(\mathbb{F}_r, [E])$,

3. $\beta \in \mathrm{Hom}_{isom}(\mathbb{F}_r, [E])$.

Observe that because $\alpha \in \mathrm{Hom}'_g(\mathbb{F}_r, [E])$ and $\epsilon > 0$ are arbitrary, this claim implies the theorem.

To see the first statement, note that

$$\mathbf{d}_u(\alpha, \beta) = d_u(\alpha(s_r), \psi) \leq \mu(B) + \mu(\cup_{i=N}^{\infty} C_i) \leq 2\epsilon.$$

For the second statement, note that the equivalence relation generated by $\{\beta(s_1), \ldots, \beta(s_r)\}$ contains the graph of $\alpha(s_1), \ldots, \alpha(s_{r-1})$ and $\alpha(s_r)|(X \setminus B)$. Because $\{\alpha(s_1), \ldots, \alpha(s_{r-1}), \alpha(s_r)|_{X \setminus B}\}$ is a graphing of E it follows that β is a graphing of E. So $\beta \in \mathrm{Hom}'_g(\mathbb{F}_r, [E])$.

For $k \geq 1$, let A_k be the set of all subgroups $H \in S(\mathbb{F}_r)$ such that $s_r^k \in H$ and if $1 \leq j < k$ then $s_r^j \notin H$. Observe that $P_i = (\mathrm{Stab}^{\beta})^{-1}(A_{i+N})$ for every $i \geq 1$. Because $B = \cup_{i=1}^{\infty} P_i$, it follows that B and γ are contained in $(\mathrm{Stab}^{\beta})^{-1}(\mathcal{F})$ where $\mathcal{F}$ is the Borel sigma-algebra of $S(\mathbb{F}_r)$. Because γ is generating for the action of $\alpha(s_1)_B$, it follows that $(\mathrm{Stab}^{\beta})^{-1}(\mathcal{F})$ contains every Borel subset of B (modulo sets of measure zero). Because $\alpha(s_1)$ is ergodic and $\mu(B) > 0$, this implies that $(\mathrm{Stab}^{\beta})^{-1}(\mathcal{F})$ is the sigma-algebra of all measurable sets of X, (up to sets of measure zero). Therefore, $\beta \in \mathrm{Hom}_{isom}(\mathbb{F}_r, [E])$ as claimed. $\square$

The following is an immediate consequence.

Corollary 4.5. *If G is a free group of rank r and (X, μ, E) is an ergodic aperiodic discrete pmp equivalence relation with cost(E) $< r$ then there exists an invariant measure $\lambda \in M(G)$ such that $(S(G), \lambda, E_G)$ is isomorphic to (X, μ, E).*

5 Encoding via sub-actions

Theorem 5.1. *Let $\mathbb{F}_r = \langle s_1, \ldots, s_r \rangle$ be the free group of rank $r \geq 2$, $\mathbb{F}'_r < \mathbb{F}_r$ be the (infinite-index) subgroup generated by $\langle s_1^2, s_2, \ldots, s_r \rangle$, K be a finite set, $\mathbb{F}_r \curvearrowright K^{\mathbb{F}_r}$ be the usual action $gx(f) := x(g^{-1}f)$. Let $X \subset K^{\mathbb{F}_r}$ be a closed invariant subspace. Then there exist subspaces $Z \subset Y \subset S(\mathbb{F}_r)$ such that*

1. *Z is $\mathbb{F}'_r$-invariant, Y is $\mathbb{F}_r$-invariant,*
2. *there is a finite set $L \subset \mathbb{F}_r$ (depending only on K) such that $Y = \bigcup_{f \in L} fZ$,*
3. *the action $\mathbb{F}_r \curvearrowright X$ is topologically conjugate to the conjugation-action $\mathbb{F}'_r \curvearrowright Z$. More precisely, if $\phi : \mathbb{F}_r \to \mathbb{F}'_r$ is the isomorphism determined by $\phi(s_1) = s_1^2, \phi(s_i) = s_i$ for $2 \leq i \leq r$ then there is a homeomorphism $\Psi : X \to Z$ such that $\Psi(fx) = \phi(f) \cdot \Psi(x)$ for all $x \in X$ and $f \in \mathbb{F}_r$.*
4. *If η is an $\mathbb{F}_r$-invariant Borel measure on X then there exists an $\mathbb{F}_r$-invariant Borel measure λ on Y such that $\Psi_*\eta = \lambda|Z$ (in particular, $\lambda(Z) > 0$). Moreover, if η is finite then λ is also finite.*

Proof. Without loss of generality, $K = \{1, \ldots, n\}$ for some integer $n \geq 1$. Given $x \in K^{\mathbb{F}_r}$, we will define a subgroup $\Psi(x) < \mathbb{F}_r$. The easiest way to understand $\Psi(x)$ is through its Schreier coset graph $\Psi(x)\backslash\mathbb{F}_r$ which is constructed as follows. Start with the Cayley graph of $\mathbb{F}_r$. For every element $g \in \mathbb{F}_r$, subdivide the edge (g, gs_1) and attach a cycle of length $x(g)$ to the new vertex. Every edge of this cycle is labeled s_2. For every $3 \leq i \leq r$, place a loop labeled s_i at all the new vertices. Also, place a loop labeled s_1 at all the new vertices other than the one subdividing the edge (g, gs_1). An example of x and $\Psi(x)\backslash G$ for $r = 2$ is shown in figure 4.

More formally, $\Psi(x)$ is the subgroup generated by all elements of the form

1. $gs_1s_2^ts_1^{-1}g^{-1}$ if $g \in \mathbb{F}'_r$ and $x(\phi^{-1}(g)) = 1$ (for any $t \in \mathbb{Z}$);
2. $gs_1s_2^ts_i^{\pm 1}s_2^us_1^{-1}g^{-1}$ if $g \in \mathbb{F}'_r$ and $x(\phi^{-1}(g)) = k > 1$ where $i \neq 2$, $t + u = 0 \mod k$, and $u, t \neq 0 \mod k$;
3. $gs_1s_is_1^{-1}g^{-1}$ if $g \in \mathbb{F}'_r$ and $3 \leq i \leq r$.



Figure 4: The element x is depicted on the left as a function on the vertices of the Cayley graph of $\mathbb{F}_2$. The Schreier coset graph $\Psi(x)\backslash G$ is depicted on the right. Black arrows represent s_1 and white arrows represent s_2.

Clearly, $\Psi(fx) = \phi(f) \cdot \Psi(x)$ for all $x \in X$ and $f \in \mathbb{F}_r$. Note $\Psi : K^{\mathbb{F}_r} \to S(\mathbb{F}_r)$ is a homeomorphism onto its image. Let $Z = \Psi(X)$, so that Ψ gives a homeomorphism from X to Z and let Y be the union of all $\mathbb{F}_r$-orbits in Z. From the construction, it is clear that if L is the radius n ball centered at the identity (with respect to the word metric) then $Y = \cup_{f \in L} f \cdot Z$. This, and the fact that Z is closed, implies Y is closed.

Now suppose η is an $\mathbb{F}_r$-invariant Borel measure on X. Let $L = \{f_1, \ldots, f_j\}$ be an ordering of L with $f_1 = e$. Define $\Upsilon : Y \to Z$ by $\Upsilon(K) = f_i \cdot K$ where i is the smallest number such that $f_i \cdot K \in Z$. Define a measure λ on Y by

$$\lambda(E) = \int |\Upsilon^{-1}(K) \cap E| \; d\Psi_*\eta(K).$$

Clearly, $\lambda|Z = \Psi_*\eta$ and λ is $\mathbb{F}_r$-invariant if and only if η is $\mathbb{F}_r$-invariant. Moreover, since Υ is finite-to-1, λ is finite if and only if η is finite. ☐

References

[ABBGNRS11] M. Abert, N. Bergeron, I. Biringer, T. Gelander, N. Nikolov, J. Raimbault and I. Samet, *On the growth of Betti numbers of locally symmetric spaces.* arXiv:1104.55

[AGV12] M. Abert, Y. Glasner and B. Virag, *Kesten's theorem for Invariant Random Subgroups.* arXiv:1201.3399

[Bo10] L. Bowen, *Random walks on coset spaces with applications to Furstenberg entropy.* arXiv:1008.4933.

[BS06] U. Bader, Y. Shalom, *Factor and normal subgroup theorems for lattices in products of groups.* Invent. Math. 163 (2006), 415-454

[CGP07] Y. de Cornulier, L. Guyot, and W. Pitsch, *On the isolated points in the space of groups.* J. Algebra 307 (2007), no. 1, 254–277.

[DS02] S. G. Dani, *On conjugacy classes of closed subgroups and stabilizers of Borel actions of Lie groups.* Ergodic Theory Dynam. Systems 22 (2002), no. 6, 1697–1714.

[FM77] J. Feldman and C. C. Moore, *Ergodic equivalence relations, cohomology, and von Neumann algebras. I*, Trans. Amer. Math. Soc. 234 (1977), no. 2, 289–324.

[Fu99] A. Furman, *Orbit Equivalence rigidity*, Ann. of Math. (2) 150 (1999), no. 3, 1083–1108.

[Gr84] R. Grigorchuk, *Degrees of growth of finitely generated groups and the theory of invariant means.* Izv. Akad. Nauk SSSR Ser. Mat., 48(5):939–985, 1984.

[Gr11] R. Grigorchuk, *Some topics of dynamics of group actions on rooted trees.*, The Proceedings of the Steklov Institute of Math., v. 273 (2011), 64–175.

[GS99] V. Golodets and S. D. Sinelshchikov, *On the conjugacy and isomorphism problems for stabilizers of Lie group actions.* Ergodic Theory Dynam. Systems 19 (1999), no. 2, 391–411.

[GW97] E. Glasner and B. Weiss, *Kazhdans property T and the geometry of the collection of invariant measures*, Geometry and Functional Analysis, vol. 7 (1997), 917–935.

[Ha44] P. R. Halmos, *In General a Measure Preserving Transformation is Mixing.* Annals of Mathematics, 45, (1944), 786–792.

[Ha58] P.R. Halmos, *Lectures on Ergodic Theory.* Chelsea, 1958.

[Hj06] G. Hjorth. *A Lemma for cost attained.* Ann. Pure Appl. Logic, 143(1-3):87–102, 2006.

[IKT09] A. Ioana, A. S. Kechris and T. Tsankov, *Subequivalence relations and positive-definite functions.* Groups Geom. Dyn. 3 (2009), no. 4, 579–625.

[Ke10] A. S. Kechris, *Global Aspects of Ergodic Group Actions.* Mathematical Surveys and Monographs, 160, American Mathematical Society, 2010.

[KM04] A. S. Kechris and B. D. Miller, *Topics in Orbit Equivalence.* Lecture Notes in Mathematics, Springer, Vol. 1852, 2004

[KST99] A. S. Kechris, S. Solecki, and S. Todorcevic, *Borel chromatic numbers.* Adv. Math., 141(1):1–44, 1999.

[LOS78] J. Lindenstrauss, G.H. Olsen, Y. Sternfeld, *The Poulsen simplex*, Ann. Inst. Fourier (Grenoble) 28 (1978), 91–114.

[Sa11] D. Savchuk, *Schreier Graphs of Actions of Thompsons Group F on the Unit Interval and on the Cantor Set*, arXiv: 1105.4017.

[SZ94] G. Stuck and R. J. Zimmer, *Stabilizers for ergodic actions of higher rank semisimple groups*, Ann. of Math. (2) 139 (1994), no. 3, 723–747.

[Ve10] A. Vershik, *Nonfree Actions of Countable Groups and their Characters*, arXiv:1012.4604

[Ve11] A. Vershik, *Totally nonfree actions and infinite symmetric group.* arXiv:1109.3413

[Vo12] Y. Vorobets, *Notes on the Schreier graphs of the Grigorchuk group.* Contemporary Mathematics 567 (2012), 221–248.

[Zi84] Zimmer, R. *Ergodic theory and semisimple groups.* Birkhauser, Boston, 1984.